

Community Health Profile  
**Public Health District Health Indicators**  
**2004 Data**

Item	Idaho	District 1	District 2	District 3	District 4	District 5	District 6	District 7
<b>Maternal/Child</b>								
1) % unintended pregnancies	<i>not available</i>							
2) % live births with adequate prenatal care <sup>1</sup>	32.6	29.0	29.1	30.6	39.9	27.5	39.1	25.4
3) % live births with low birth weight <sup>2</sup>	6.8	6.0	5.6	6.4	7.4	6.6	7.5	6.9
4) % live births with tobacco use during pregnancy <sup>3</sup>	13.1	20.7	16.2	12.3	12.8	14.7	12.9	7.3
5) % WIC participation <sup>4</sup>	39.3	41.3	41.0	42.9	26.0	43.0	52.0	42.8
6) % prevalence of breastfeeding <sup>5</sup>	88.7	87.0	89.0	89.5	92.1	85.5	84.2	89.1
<b>Adolescents</b>								
1) Teen pregnancy rate (15-19) <sup>6</sup>	45.2	40.7	32.8	64.0	38.7	61.5	38.6	37.5
2) Motor vehicle crash death rate, 2002-2004 (ages 15-19) <sup>7,9</sup>	30.9	37.6	36.3	19.6	13.9	51.0	43.2	34.2
3) Suicide rate, 2002-2004 (ages 10-18) <sup>8,9</sup>	5.7	1.3	8.2	4.3	7.6	6.8	2.7	8.8
<b>Adults</b>								
1) % without health care coverage	15.5	19.6	15.9	18.9	11.7	16.8	14.0	14.9
2) % with no leisure time activity	18.8	19.1	16.3	27.3	14.0	24.0	17.2	16.6
3) % overweight (BMI > 25)	58.2	60.0	59.1	60.1	54.3	60.5	57.8	60.4
4) % diagnosed with diabetes	6.2	4.6	6.7	7.2	4.3	7.0	8.9	7.2
5) % who smoke cigarettes	17.4	20.9	18.6	19.7	15.6	20.0	16.6	12.2
6) % who binge drink (5+ drinks on one occasion in past 30 days)	12.8	14.1	20.6	13.6	12.1	13.9	10.0	8.1
7) % of females without breast cancer screening (age 40+) <sup>10</sup>	40.7	46.6	39.4	42.3	30.6	44.3	42.1	49.6
8) % of males without prostate cancer screening (age 40+) <sup>11</sup>	52.0	56.5	50.7	55.9	46.0	54.9	55.6	49.0
9) % who did not wear seatbelts (2003)	32.7	30.0	30.7	31.7	26.0	35.8	41.0	43.0
10) Suicide rate, 2002-2004 (ages 65+) <sup>8,9</sup>	22.5	31.3	29.4	24.2	18.5	13.7	24.5	17.9
11) % with no dental visit in the past 12 months	33.4	37.5	36.4	38.6	28.8	38.2	32.3	26.8

**SOURCES:**

For Maternal/Child indicators 2-6, the adolescent indicators, and adult indicator #10: Vital Statistics, Bureau of Health Policy and Vital Statistics  
For adults items 1-9 and 11: Idaho Behavioral Risk Factor Surveillance System, Bureau of Health Policy and Vital Statistics

**NOTES:**

1. In 2004, the Idaho birth certificate was revised. Beginning in 2004, Idaho prenatal care data are based on date of first prenatal care visit as reported in the mother's medical record. 2004 APNCU data are not comparable with APNCU data based on month prenatal care began. Percents based on records with stated data for APNCU calculation (date of first prenatal care visit, number of visits, and length of gestation).
2. Percents based on records with stated birth weight.
3. Prior to 2004, tobacco use during pregnancy (yes or no) was reported on the Idaho birth certificate. Beginning in 2004, cigarette smoking before and during pregnancy is reported on the revised birth certificate. Data in 2004 are not comparable to data prior to 2004.
4. The revised 2004 Idaho birth certificate includes a new variable for reporting if mother received WIC food for herself during this pregnancy (yes or no).
5. The revised 2004 birth certificate includes a new variable for reporting if infant was being breastfed at birth or at the time the birth certificate was completed (yes or no).
6. Pregnancies include live births, induced abortions, and reportable stillbirths (only those fetal deaths with a gestational period of 20+ weeks or which weight 350+ grams are required to be reportable under Idaho law.) Rate: Number of pregnancies per 1,000 females in corresponding age group. Rates are based on July 1, 2004 female population estimates from the National Center for Health Statistics, United States resident population from the Vintage 2004 postcensal series, prepared under a collaborative arrangement with the U.S. Census Bureau, Internet release date September 9, 2005.
7. Motor Vehicle Crash Death Rate: number of motor vehicle crash deaths from 2002 to 2004 per 100,000 population.
8. Suicide rate: number of suicides from 2002 to 2004 per 100,000 population.
9. Rates are based on the National Center for Health Statistics, estimates of the July 1, 2003 United States resident population from the Vintage 2003 postcensal series, prepared under a collaborative arrangement with the U.S. Census Bureau, Internet release date August 18, 2004.
10. Breast cancer screening: mammogram and clinical breast exam in past two years among females 40 and over.
11. Prostate cancer screening: Prostate-Specific Antigen (PSA) test in past two years among males 40 and over.
12. Seatbelt usage question was not asked in 2004. Reported figures are from BRFSS 2003.